# Overview of Current ARM Cloud Parameterization and Modeling Activities, Data, and Plans

Stephen Klein
Lawrence Livermore National Laboratory

November 3, 2004

Joint ARM CPM – CERES - GCSS WG4 Meeting Williamsburg, Virginia

**UCRL-PRES-xxxx** 



#### **Outline**

- SCM & CRM Forcing Data Sets for ARM Intensive Observing Periods (IOPs) at the Southern Great Plains site and selected results from their use
- "Continuous" Forcing Data Sets and selected results from their use
- Broad-Band Heating Rate Profile (BBHRP) project
- Future ARM IOPs and the ARM Mobile Facility




# **ARM** goals


- A broad goal of the ARM program is to improve the simulation of clouds and their related processes through the detailed analysis of ARM data and Cloud Resolving Models
- The technique we have focussed on is the simulation for selected ARM IOPs by Cloud Resolving Models (CRMs) and Single-Column-Models (SCMs) driven by ARM data



# Single Column Model (SCM)



# Cloud Resolving Model (CRM)





# The SCM/CRM Forcing Approach

"Advective Forcing" computed from observations



- CRM "explicitly"
- SCM parameterizations


$$\frac{\partial s}{\partial t} = \begin{bmatrix} -V \bullet \nabla s - \omega \frac{\partial s}{\partial p} \end{bmatrix}$$

$$\frac{\partial q}{\partial t} = -V \cdot \nabla q - \omega \frac{\partial q}{\partial p}$$

$$+L(c-e)-\frac{\partial \overline{\omega's'}}{\partial p}+Q_{rad}$$

$$(e-c)-\frac{\partial \overline{\omega'q'}}{\partial p}$$





### **ARM Data During IOPs**


- 5 sounding stations
- 7 wind profilers
- Dozens of surface flux and radiation estimates
- Surface observations from Oklahoma and Kansas Mesonet
- Precipitation estimates from radar
- TOA satellite estimates (Minnis et al.)
- + NCEP Regional Model (RUC) background fields

Variational Analysis (Zhang and Lin 1997, Zhang et al. 2001)

The profiles of state variables are adjusted until they satisfy column budgets of mass, energy, and moisture

# Forcing Example

#### Strong Frontal Passage, March 2-3, 2000





# IOPs Analyzed (and data available for...)

- Summer: 1995, 1997, 1999, 2002
- Fall: 1997, 2000, 2002
- Winter: 1999, 2000
- Spring: 1997, 1998, 1999, 2000

A total of ~250 days


#### Two IOPs have have been intensely studied

- Case 3 June-July 1997 Summer Deep Convection
- Case 4 March 2000 Spring Frontal Clouds

#### **Case 3 Results**

(Xie et al. 2002, Xu et al. 2002)

#### Case 3 mean T and q errors for 10 CRMs and 11 SCMs




CRMs produce less bias and less scatter than SCMs (they should!)

Root mean square errors are also less.



# **Case 3 Relative Humidity and Cloud Fraction**






CRMs maintain a better relative humidity and cloud fraction structure

SCMs are too moist in relative humidity and associated cloud aloft

**ARM Cloud Radar Observation** 



## SCM errors may be caused by a lack of downdrafts



Net cumulus mass flux

**Updraft mass flux** 


**Downdraft mass flux** 



### **Case 4 Results**

(Xie et al. 2005, Xu et al. 2005)

#### Strong Frontal Passage – March 2-3, 2000



Cloud Frequency (%)

165
265
365
465
565
665
765
865
965
18 00 06 12 18 00 06 12 18 00
MAR 2 MAR3
0 20 40 60 80 100
10 30 50 70 90


IR Image, 12Z March 2

**ARM Cloud Radar Observation** 




#### **Cloud Fraction**

#### **Observation**



**Selected CRMs** 





CRM clouds miss low level clouds

CRMs are too "convective"

**Selected SCMs** 


SCM clouds are too extensive





### **Cloud Liquid**

#### **Observation**




'Microbase' analysis courtesy of Mark Miller et al.

**Selected CRMs** 

**Selected SCMs** 

SCM liquid water varies a lot between models



#### **Cloud Ice**

#### **Observation**

165 - ice (b)

265 - 465 - 665 - 765 - 865 - 865 - 965 - 18 00 06 12 18 00 06 12 18 00 MAR 2 MAR 3


'Microbase' analysis courtesy of Mark Miller et al.

#### **Selected CRMs**

CRM ice water varies a lot between models

**Selected SCMs** 

SCM ice water varies a lot between models



# "Continuous" Forcing

(Xie et al. 2004)

The Variational Analysis forcing is so nice, wouldn't it be great if it weren't limited to ARM IOPs?

Yes!


If we do this, what observations would you miss?


Primarily the radiosondes... but it is still worth
doing (and produces forcing which is superior to
NWP output – at least in summer).

Three years have been completed – 1999 - 2001



# June 2000 "Continuous" Forcing





# **Examples of "Continuous" Forcing Uses**

Build statistics! Move beyond case studies!

From the JGR Special Issue of the ARM CPM Working Group:

Gordon et al. 2005, "Cluster analysis of cloud regimes and characteristic dynamics of midlatitude synoptic systems in observations and a model"


Del Genio et al. 2005, "Evaluation of regional cloud feedbacks with Single Column Models"

**GISS SCM** 

500 mb Ascent

ISCCP Cloud Radar .3 3.6 9.4 23 6 CLOUD OPTICAL THICKNESS CLOUD PHYSICAL THICKNESS (KM) SCM MEAN SCM MEAN OMEGA < 0 OMEGA < 0 CLOUD OPTICAL THICKNESS CLOUD PHYSICAL THICKNESS(KM)

500 mb Descent



# Broad-Band Heating Rate Profile Project (BBHRP)

Eli Mlawer, Mark Miller, and many others, ...

"So the first question that we need to answer is: Given a specified three-dimensional field of cloud properties, can we compute with sufficient accuracy the solar and terrestrial radiative flux transfer and associated atmospheric heating rates through the clouds?"

Ackerman and Stokes, Physics Today

- The goal is to derive the vertical profile of radiative heating rates over the ARM site continuously given all of ARM measurements and TOA satellite fluxes
- First do locally to the Central Facility, then extend to the whole ARM SGP site
- Uses the best ARM measurements of aerosol optical depth, LWP, etc.
- Radar clouds and the retrieved cloud micropohysics are key
- Focussed on the period 1999-2001
- First at central facility, then whole SGP domain, then other ARM sites




# **ARM Field Projects**

Mixed Phase Arctic Cloud Experiment (MPACE)

North Slope of Alaska October 2004

**♦** see upcoming talk by Hans Verlinde



Tropical Warm Pool – International Cloud Experiment (TWP-ICE)

Darwin Australia

January – February 2006

♦ see upcoming talk by Jim Mather





# **ARM Mobile Facility Deployments**



Marine Stratus Radiation Aerosol and Drizzle (MASRAD)

California Coast


March – September 2005



RAdiative Divergence using AMF, GERB, and AMMA STations (RADAGAST)

Niamey, Niger

~ January - December 2006





# The End



# Case 3 RMS T, q, and RH errors

